

# i-Tree Ecosystem Analysis

## City of Pacific Grove



Urban Forest Effects and Values  
June 2011

---



## Summary

Understanding an urban forest's structure, function and value can promote management decisions that will improve human health and environmental quality. An assessment of the vegetation structure, function, and value of the City of Pacific Grove urban forest was conducted during 2010. Data from collected 126 field plots (out of total 157 plots) located throughout City of Pacific Grove were analyzed using the Urban Forest Effects (UFORE) model developed by the U.S. Forest Service, Northern Research Station.

### Key findings

- Number of trees: 25,900
- Tree cover: 23.1%
- Most common species: Monterey pine, Coastal live oak, Monterey cypress
- Percentage of trees less than 6" (15.2 cm) diameter: 11.3%
- Pollution removal: 23 metric tons/year (\$177 thousand/year)
- Carbon storage: 10,900 metric tons (\$222 thousand)
- Carbon sequestration: 890 metric tons/year (\$18.1 thousand/year)
- Building energy savings: \$12.4 thousand / year
- Avoided carbon emissions: \$-81 / year
- Structural values: \$209 million

Metric ton: 1000 kilograms

Carbon storage: the amount of carbon bound up in the above-ground and below-ground parts of woody vegetation

Carbon sequestration: the removal of carbon dioxide from the air by plants through photosynthesis

Structural value: value based on the physical resource itself (e.g., the cost of having to replace a tree with a similar tree)

For an overview of UFORE methodology, see Appendix I. Data collection quality is determined by the local data collectors, over which i-Tree has no control. Additionally, some of the plot and tree information may not have been collected, so not all of the analyses may have been conducted for this report.

## Table of Contents

Summary.....	2
I. Tree characteristics of the urban forest.....	4
II. Urban forest cover & leaf area.....	6
III. Air pollution removal by urban trees.....	7
IV. Carbon storage and sequestration.....	8
V. Trees and building energy use.....	9
VI. Structural and functional values.....	10
VII. Potential pest impacts.....	11
Appendix I. UFORE Model and Field Measurements.....	12
Appendix II. Relative tree effects.....	13
Appendix III. Comparison of urban forests.....	14
Appendix IV. General recommendations for air quality improvement..	15
References.....	16

# I. Tree Characteristics of the Urban Forest

The urban forest of City of Pacific Grove has an estimated 25,900 trees with a tree cover of 23.1 percent. Trees that have diameters less than 6-inches constitute 11.3 percent of the population. The three most common species are Monterey pine (34.30 percent), Coastal live oak (28.80 percent), and Monterey cypress (21.20 percent).




Figure 1. Tree species composition in City of Pacific Grove

Among the land use categories, the highest tree densities occur in ForestGrove\_DMP followed by PG\_Retreat and SunsetDrive. The overall tree density in City of Pacific Grove is 35.8 trees / hectare (see Appendix III for comparable values from other cities).






Figure 2. Number of trees/ha in City of Pacific Grove by land use



**Figure 3. Percent of tree population by diameter class (DBH=stem diameter at 1.37 meter)**

Urban forests are composed of a mix of native and exotic tree species. Thus, urban forests often have a tree diversity that is higher than surrounding native landscapes. An increased tree diversity can minimize the overall impact or destruction by a species-specific insect or disease, but it can also pose a risk to native plants if some of the exotic species are invasive plants that can potentially out-compete and displace native species. In City of Pacific Grove, about 86 percent of the trees are from species native to North America, while 85 percent are native to the state or district. Species exotic to California make up 14 percent of the population. Most exotic tree species have an origin from Australia (4.4 percent of the species).



**Figure 4. Percent of live trees by species origin**

"North America +" = native to North America and at least one other continent except South America

"Americas +" = native to North and South America and at least one other continent

## II. Urban Forest Cover and Leaf Area


Many tree benefits equate directly to the amount of healthy leaf surface area of the plant. In City of Pacific Grove, the three most dominant species in terms of leaf area are Monterey pine, Monterey cypress, and Coastal live oak. Trees cover about 23.1 percent of City of Pacific Grove, and shrubs cover 13.2 percent.

The 10 most important species are listed in the table below. Importance values (IV) are calculated as the sum of relative leaf area and relative composition.

**Table 1. Most important species in City of Pacific Grove**

Common Name	Percent Population	Percent Leaf Area	IV
Monterey pine	34.3	42.7	77.0
Monterey cypress	21.2	31.6	52.8
Coastal live oak	28.8	12.4	41.2
Coast redwood	0.6	3.2	3.7
Black acacia	1.8	1.7	3.6
Red gum eucalyptus	1.1	2.0	3.2
Cladrastis spp	0.5	1.3	1.8
Willow spp	0.7	1.1	1.8
Victorian box	1.1	0.2	1.2
Flowering plum	1.1	0.1	1.2

The two most dominant ground cover types are Tar (26.6 percent) and Building (20.4 percent).



**Figure 5. Percent ground cover in City of Pacific Grove**

### III. Air Pollution Removal by Urban Trees

Poor air quality is a common problem in many urban areas. It can lead to decreased human health, damage to landscape materials and ecosystem processes, and reduced visibility. The urban forest can help improve air quality by reducing air temperature, directly removing pollutants from the air, and reducing energy consumption in buildings, which consequently reduces air pollutant emissions from the power plants. Trees also emit volatile organic compounds that can contribute to ozone formation. However, integrative studies have revealed that an increase in tree cover leads to reduced ozone formation[1].

Pollution removal by trees and shrubs in City of Pacific Grove was estimated using field data and recent pollution and weather data available. Pollution removal was greatest for ozone. It is estimated that trees and shrubs remove 23 metric tons of air pollution (ozone (O3), carbon monoxide (CO), nitrogen dioxide (NO2), particulate matter less than 10 microns (PM10), and sulfur dioxide (SO2)) per year with an associated value of \$177 thousand (based on estimated national median externality costs associated with pollutants[2]).





Figure 6. Pollution removal and associated value for trees in City of Pacific Grove (line graph is value)

## IV. Carbon Storage and Sequestration

Climate change is an issue of global concern. Urban trees can help mitigate climate change by sequestering atmospheric carbon (from carbon dioxide) in tissue and by altering energy use in buildings, and consequently altering carbon dioxide emissions from fossil-fuel based power plants[3].

Trees reduce the amount of carbon in the atmosphere by sequestering carbon in new growth every year. The amount of carbon annually sequestered is increased with the size and health of the trees. The gross sequestration of City of Pacific Grove trees is about 890 metric tons of carbon per year with an associated value of \$18.1 thousand. Net carbon sequestration in the urban forest is about 819 metric tons.



**Figure 7. Carbon sequestration and value for species with greatest overall carbon sequestration in City of Pacific Grove**

As trees grow they store more carbon as wood. As trees die and decay, they release much of the stored carbon back to the atmosphere. Thus, carbon storage is an indication of the amount of carbon that can be lost if trees are allowed to die and decompose. Trees in City of Pacific Grove are estimated to store 10,900 metric tons of carbon (\$222 thousand). Of the species sampled, Coastal live oak stores and sequesters the most carbon (approximately 36.6% of the total carbon stored and 45.9% of all sequestered carbon.)

## V. Trees and Building Energy Use

Trees affect energy consumption by shading buildings, providing evaporative cooling, and blocking winter winds. Trees tend to reduce building energy consumption in the summer months and can either increase or decrease building energy use in the winter months, depending on the location of trees around the building. Estimates of tree effects on energy use are based on field measurements of tree distance and direction to space conditioned residential buildings[4].

Based on 2002 prices, trees in City of Pacific Grove are estimated to reduce energy-related costs from residential buildings by \$12.4 thousand annually. Trees also provide an additional \$-81 in value[5] by reducing the amount of carbon released by fossil-fuel based power plants (a reduction of -4 metric tons of carbon emissions).

**Table 2. Annual energy savings due to trees near residential buildings. Note: negative numbers indicate an increased energy use or carbon emission.**

	Heating	Cooling	Total
MBTU <sup>1</sup>	-1,226	n/a	-1,226
MWH <sup>2</sup>	-61	221	160
Carbon avoided (mt)	-25	21	-4

<sup>1</sup>One million British Thermal Units

<sup>2</sup>Megawatt-hour

**Table 3. Annual savings<sup>1</sup> (US \$) in residential energy expenditure during heating and cooling seasons. Note: negative numbers indicate a cost due to increased energy use or carbon emission.**

	Heating	Cooling	Total
MBTU <sup>2</sup>	-8,486	n/a	-8,486
MWH <sup>3</sup>	-7,950	28,804	20,854
Carbon avoided (mt)	-507	426	-81

<sup>1</sup>Based on state-wide energy costs for California.

<sup>2</sup>One million British Thermal Units

<sup>3</sup>Megawatt-hour

## VI. Structural and Functional Values

Urban forests have a structural value based on the trees themselves (e.g., the cost of having to replace a tree with a similar tree); they also have functional values (either positive or negative) based on the functions the trees perform.

The structural value of an urban forest tends to increase with a rise in the number and size of healthy trees [6]. Annual functional values also tend to increase with increased number and size of healthy trees, and are usually on the order of several million dollars per year. Through proper management, urban forest values can be increased; however, the values and benefits also can decrease as the amount of healthy tree cover declines.

### Structural values:

- Structural value: \$209 million
- Carbon storage: \$222 thousand

### Annual functional values:

- Carbon sequestration: \$18.1 thousand
- Pollution removal: \$177 thousand
- Lower energy costs and carbon emission reductions: \$12.3 thousand (Note: negative value indicates increased energy cost and carbon emission value)





Figure 8. Structural value of the 10 most valuable tree species in City of Pacific Grove

## VII. Potential Pest Impacts

Various insects and diseases can infest urban forests, potentially killing trees and reducing the health, value and sustainability of the urban forest. As pests tend to have differing tree hosts, the potential damage or risk of each pest will differ. Four exotic pests were analyzed for their potential impact: Asian longhorned beetle (ALB), gypsy moth (GM), emerald ash borer (EAB), and Dutch elm disease (DED).



**Figure 9. Number of susceptible City of Pacific Grove trees and structural value by pest (line graph is structural value)**

The Asian longhorned beetle (ALB) [7] is an insect that bores into and kills a wide range of hardwood species. ALB poses a threat to 4 percent of the City of Pacific Grove urban forest, which represents a loss of \$7.69 million in damage to the structure.

The gypsy moth (GM)[8] is a defoliator that feeds on many species causing widespread defoliation and tree death if outbreak conditions last several years. This pest threatens 30.6 percent of the population, which represents a loss of \$59.4 million in structural value.

Emerald ash borer (EAB)[9] has killed thousands of ash trees in parts of the United States. EAB has the potential to affect 0 percent of the population (\$0 in structural damage).

American elm, one of the most important street trees in the twentieth century, has been devastated by the Dutch elm disease (DED)[10]. Since first reported in the 1930s, it has killed over 50 percent of the native elm population in the United States. Although some elm species have shown varying degrees of resistance, City of Pacific Grove could possibly lose 0 percent of its trees to this pest (\$0 in structural value).

## Appendix I. UFORE Model and Field Measurements

UFORE is designed to use standardized field data from randomly located plots and local hourly air pollution and meteorological data to quantify urban forest structure and its numerous effects [5], including:

- Urban forest structure (e.g., species composition, tree health, leaf area, etc.).
- Amount of pollution removed hourly by the urban forest, and its associated percent air quality improvement throughout a year. Pollution removal is calculated for ozone, sulfur dioxide, nitrogen dioxide, carbon monoxide and particulate matter (<10 microns).
- Total carbon stored and net carbon annually sequestered by the urban forest.
- Effects of trees on building energy use and consequent effects on carbon dioxide emissions from power plants.
- Structural value of the forest, as well as the value for air pollution removal and carbon storage and sequestration.
- Potential impact of infestations by Asian longhorned beetles, emerald ash borers, gypsy moth, and Dutch elm disease.

In the field 0.040 hectare plots were randomly distributed. Typically, all field data are collected during the leaf-on season to properly assess tree canopies. Within each plot, typical data collection (actual data collection may vary depending upon the user) includes land use, ground and tree cover, individual tree attributes of species, stem diameter, height, crown width, crown canopy missing and dieback, and distance and direction to residential buildings[11].

To calculate current carbon storage, biomass for each tree was calculated using equations from the literature and measured tree data. Open-grown, maintained trees tend to have less biomass than predicted by forest-derived biomass equations[12]. To adjust for this difference, biomass results for open-grown urban trees were multiplied by 0.8. No adjustment was made for trees found in natural stand conditions. Tree dry-weight biomass was converted to stored carbon by multiplying by 0.5.

To estimate the gross amount of carbon sequestered annually, average diameter growth from the appropriate genera and diameter class and tree condition was added to the existing tree diameter (year  $x$ ) to estimate tree diameter and carbon storage in year  $x+1$ .

Air pollution removal estimates are derived from calculated hourly tree-canopy resistances for ozone, and sulfur and nitrogen dioxides based on a hybrid of big-leaf and multi-layer canopy deposition models[13,14]. As the removal of carbon monoxide and particulate matter by vegetation is not directly related to transpiration, removal rates (deposition velocities) for these pollutants were based on average measured values from the literature[15,16] that were adjusted depending on leaf phenology and leaf area. Particulate removal incorporated a 50 percent resuspension rate of particles back to the atmosphere[17].

If appropriate field data were collected, seasonal effects of trees on residential building energy use were calculated based on procedures described the literature[4] using distance and direction of trees from residential structures, tree height and tree condition data.

Structural values were based on valuation procedures of the Council of Tree and Landscape Appraisers[8], which uses tree species, diameter, condition and location information[18].

## Appendix II. Relative Tree Effects

The urban forest in City of Pacific Grove provides benefits that include carbon storage and sequestration, and air pollutant removal. To estimate the relative value of these benefits, tree benefits were compared to estimates of average municipal carbon emissions[19], average passenger automobile emissions[20], and average household emissions[21].

### Carbon storage is equivalent to:

- Amount of carbon emitted in City of Pacific Grove in 15 days
- Annual carbon (C) emissions from 7,240 automobiles
- Annual C emissions from 3,630 single-family houses

### Carbon monoxide removal is equivalent to:

- Annual carbon monoxide emissions from 6 automobiles
- Annual carbon monoxide emissions from 25 single-family houses

### Nitrogen dioxide removal is equivalent to:

- Annual nitrogen dioxide emissions from 304 automobiles
- Annual nitrogen dioxide emissions from 203 single-family houses

### Sulfur dioxide removal is equivalent to:

- Annual sulfur dioxide emissions from 2,070 automobiles
- Annual sulfur dioxide emissions from 35 single-family houses

### Particulate matter less than 10 micron (PM10) removal is equivalent to:

- Annual PM10 emissions from 22,200 automobiles
- Annual PM10 emissions from 2,150 single-family houses

### Annual carbon sequestration is equivalent to:

- Amount of carbon emitted in City of Pacific Grove in 1.2 days
- Annual C emissions from 600 automobiles
- Annual C emissions from 300 single-family houses

Note: estimates above are partially based on the user-supplied information on human population total for study area

## Appendix III. Comparison of Urban Forests

A common question asked is, "How does this city compare to other cities?" Although comparison among cities should be made with caution as there are many attributes of a city that affect urban forest structure and functions, summary data are provided from other cities analyzed using the UFORE model.

### I. City totals for trees

City	% Tree Cover	Number of trees	Carbon storage (metric tons)	Carbon Sequestration (metric tons/yr)	Pollution removal (metric tons/yr)	Pollution Value (\$US)
Calgary, Canada	7.2	11,889,000	404,000	19,400	296	1,611,000
Atlanta, GA	36.8	9,415,000	1,220,000	42,100	1,508	2,534,000
Toronto, Canada	20.5	7,542,000	900,000	36,600	1,100	6,105,000
New York, NY	21	5,212,000	1,226,000	38,400	1,521	8,071,000
Baltimore, MD	21	2,627,000	541,000	14,600	390	2,129,000
Philadelphia, PA	15.7	2,113,000	481,000	14,600	523	2,826,000
Washington, DC	28.6	1,928,000	474,000	14,600	379	1,956,000
Boston, MA	22.3	1,183,000	289,000	9,500	258	1,426,000
Woodbridge, NJ	29.5	986,000	145,000	5,000	191	1,037,000
Minneapolis, MN	26.5	979,000	227,000	8,100	277	1,527,000
Syracuse, NY	23.1	876,000	157,000	4,900	99	268,000
Morgantown, WV	35.9	661,000	85,000	2,700	60	311,000
Moorestown, NJ	28	583,000	106,000	3,400	107	576,000
Jersey City, NJ	11.5	136,000	19,000	800	37	196,000
Freehold, NJ	34.4	48,000	18,000	500	19	133,000

### II. Per hectare values of tree effects

City	No. of trees	Carbon Storage (metric tons)	Carbon sequestration (kgs/yr)	Pollution removal (kgs/yr)	Pollution Value (\$US)
Calgary, Canada	164.8	5.60	0.13	4.0	22.2
Atlanta, GA	275.8	35.64	0.62	44.2	74.1
Toronto, Canada	119.4	14.35	0.29	17.5	96.6
New York, NY	65.2	15.24	0.24	19.1	101.1
Baltimore, MD	125.5	25.78	0.35	18.6	101.8
Philadelphia, PA	61.8	14.12	0.21	15.2	82.8
Washington, DC	121.1	29.81	0.46	23.8	122.8
Boston, MA	82.8	20.18	0.33	17.9	99.8
Woodbridge, NJ	164.3	24.21	0.42	31.8	173.0
Minneapolis, MN	64.7	15.02	0.27	18.4	101.1
Syracuse, NY	134.7	24.21	0.38	15.2	41.3
Morgantown, WV	295.8	38.11	0.60	26.7	139.1
Moorestown, NJ	153.2	28.02	0.45	28.2	151.5
Jersey City, NJ	35.3	4.93	0.11	9.6	51.2
Freehold, NJ	95.1	35.87	0.49	37.7	263.4

## Appendix IV. General Recommendations for Air Quality Improvement

Urban vegetation can directly and indirectly affect local and regional air quality by altering the urban atmosphere environment. Four main ways that urban trees affect air quality are[22]:

- Temperature reduction and other microclimate effects
- Removal of air pollutants
- Emission of volatile organic compounds (VOC) and tree maintenance emissions
- Energy effects on buildings

The cumulative and interactive effects of trees on climate, pollution removal, and VOC and power plant emissions determine the impact of trees on air pollution. Cumulative studies involving urban tree impacts on ozone have revealed that increased urban canopy cover, particularly with low VOC emitting species, leads to reduced ozone concentrations in cities[23]. Local urban management decisions also can help improve air quality.

Urban forest management strategies to help improve air quality include[24]:

Strategy	Result
Increase the number of healthy trees	Increase pollution removal
Sustain existing tree cover	Maintain pollution removal levels
Maximize use of low VOC-emitting trees	Reduces ozone and carbon monoxide formation
Sustain large, healthy trees	Large trees have greatest per-tree effects
Use long-lived trees	Reduce long-term pollutant emissions from planting and removal
Use low maintenance trees	Reduce pollutants emissions from maintenance activities
Reduce fossil fuel use in maintaining vegetation	Reduce pollutant emissions
Plant trees in energy conserving locations	Reduce pollutant emissions from power plants
Plant trees to shade parked cars	Reduce vehicular VOC emissions
Supply ample water to vegetation	Enhance pollution removal and temperature reduction
Plant trees in polluted or heavily populated areas	Maximizes tree air quality benefits
Avoid pollutant-sensitive species	Improve tree health
Utilize evergreen trees for particulate matter	Year-round removal of particles

## References

1. Nowak D.J. and Dwyer J.F. "Understanding the Benefits and Costs of Urban Forest Ecosystems." Handbook of Urban and Community Forestry in the Northeast. Ed. John E. Kuser. Kluwer Academics/Plenum Pub., New York. 2000. 11-22.
2. Murray, F.J.; Marsh L.; Bradford, P.A. 1994. New York State Energy Plan, vol. II: issue reports. Albany, NY: New York State Energy Office.
3. Abdollahi, K.K.; Z.H. Ning; and A. Appeaning (eds). 2000. Global climate change and the urban forest. Baton Rouge, LA: GCRCC and Franklin Press. 77p.
4. McPherson, E.G. and J. R. Simpson 1999. Carbon dioxide reduction through urban forestry: guidelines for professional and volunteer tree planters. Gen. Tech. Rep. PSW-171. Albany, CA: U.S. Department of Agriculture, Forest Service, Pacific Southwest Research station 237 p. [http://wcufrre.ucdavis.edu/products/cufr\\_43.pdf](http://wcufrre.ucdavis.edu/products/cufr_43.pdf)
5. Nowak, D.J., and D.E. Crane. 2000. The Urban Forest Effects (UFORE) Model: quantifying urban forest structure and functions. In: Hansen, M. and T. Burk (Eds.) Integrated Tools for Natural Resources Inventories in the 21st Century. Proc. Of the IUFRO Conference. USDA Forest Service General Technical Report NC-212. North Central Research Station, St. Paul, MN. pp. 714-720. See also <http://www.ufore.org>.
6. Nowak, D.J.; Crane, D.E.; Dwyer, J.F. 2002. Compensatory value of urban trees in the United States. Journal of Arboriculture. 28(4): 194 - 199.
7. Northeastern Area State and Private Forestry. 2005. Asian Longhorned Beetle. Newtown Square, PA: U.S. Department of Agriculture, Northeastern Area State and Private Forestry. <http://www.na.fs.fed.us/spfo/alb/>
8. Northeastern Area State and Private Forestry. 2005. Gypsy moth digest. Newtown Square, PA: U.S. Department of Agriculture, Forest Service, Northeastern Area State and Private Forestry. <http://na.fs.fed.us/fhp/gm>
9. Northeastern Area State and Private Forestry. 2005. Forest health protection emerald ash borer home. Newtown Square, PA: U.S. Department of Agriculture, Forest Service, Northeastern Area State and Private Forestry. <http://www.na.fs.fed.us/spfo/eab/index.html>
10. Northeastern Area State and Private Forestry. 1998. HOW to identify and manage Dutch Elm Disease. NA-PR-07-98. Newtown Square, PA: U.S. Department of Agriculture, Forest Service, Northeastern Area State and Private Forestry. [http://www.na.fs.fed.us/spfo/pubs/howtos/ht\\_ded/ht\\_ded.htm](http://www.na.fs.fed.us/spfo/pubs/howtos/ht_ded/ht_ded.htm)
11. Nowak, D.J.; Crane, D.E.; Stevens, J.C.; Hoehn, R.E. 2005. The urban forest effects (UFORE) model: field data collection manual. V1b. Newtown Square, PA: U.S. Department of Agriculture, Forest Service, Northeastern Research Station, 34 p. [http://www.fs.fed.us/ne/syracuse/Tools/downloads/UFORE\\_Manual.pdf](http://www.fs.fed.us/ne/syracuse/Tools/downloads/UFORE_Manual.pdf)

12. Nowak, D.J. 1994. Atmospheric carbon dioxide reduction by Chicago's urban forest. In: McPherson, E.G.; Nowak, D.J.; Rowntree, R.A., eds. Chicago's urban forest ecosystem: results of the Chicago Urban Forest Climate Project. Gen. Tech. Rep. NE-186. Radnor, PA: U.S. Department of Agriculture, Forest Service, Northeastern Forest Experiment Station: 83-94.

13. Baldocchi, D. 1988. A multi-layer model for estimating sulfur dioxide deposition to a deciduous oak forest canopy. *Atmospheric Environment*. 22: 869-884.

14. Baldocchi, D.D.; Hicks, B.B.; Camara, P. 1987. A canopy stomatal resistance model for gaseous deposition to vegetated surfaces. *Atmospheric Environment*. 21: 91-101.

15. Bidwell, R.G.S.; Fraser, D.E. 1972. Carbon monoxide uptake and metabolism by leaves. *Canadian Journal of Botany*. 50: 1435-1439.

16. Lovett, G.M. 1994. Atmospheric deposition of nutrients and pollutants in North America: an ecological perspective. *Ecological Applications*. 4: 629-650.

17. Zinke, P.J. 1967. Forest interception studies in the United States. In: Sopper, W.E.; Lull, H.W., eds. *Forest Hydrology*. Oxford, UK: Pergamon Press: 137-161.

18. Nowak, D.J.; Crane, D.E.; Stevens, J.C.; Ibarra, M. 2002. Brooklyn's Urban Forest. Gen. Tech. Rep. NE-290. Newtown Square, PA: U.S. Department of Agriculture, Forest Service, Northeastern Research Station. 107 p. Council of Tree and Landscape Appraisers guidelines. For more information, see Nowak, D.J., D.E. Crane, and J.F. Dwyer. 2002. Compensatory value of urban trees in the United States. *J. Arboric.* 28(4): 194-199.

19. Total city carbon emissions were based on 2003 U.S. per capita carbon emissions - calculated as total U.S. carbon emissions (Energy Information Administration, 2003, *Emissions of Greenhouse Gases in the United States 2003*. <http://www.eia.doe.gov/oiaf/1605/ggrpt/>) divided by 2003 U.S. total population ([www.census.gov](http://www.census.gov)). Per capita emissions were multiplied by city population to estimate total city carbon emissions.

20. Average passenger automobile emissions per mile were based on dividing total 2002 pollutant emissions from light-duty gas vehicles (National Emission Trends <http://www.epa.gov/ttn/chief/trends/index.html>) divided by total miles driven in 2002 by passenger cars (National Transportation Statistics [http://www.bts.gov/publications/national\\_transportation\\_statistics/2004/](http://www.bts.gov/publications/national_transportation_statistics/2004/)).

Average annual passenger automobile emissions per vehicle were based on dividing total 2002 pollutant emissions from light-duty gas vehicles by total number of passenger cars in 2002 (National Transportation Statistics [http://www.bts.gov/publications/national\\_transportation\\_statistics/2004/](http://www.bts.gov/publications/national_transportation_statistics/2004/)).

Carbon dioxide emissions from automobile assumed six pounds of carbon per gallon of gasoline if energy costs of refinement and transportation are included (Graham, R.L., Wright, L.L., and Turhollow, A.F. 1992. The potential for short-rotation woody crops to reduce U.S. CO<sub>2</sub> Emissions. *Climatic Change* 22:223-238.

21. Average household emissions based on average electricity kWh usage, natural gas Btu usage, fuel oil Btu usage, kerosene Btu usage, LPG Btu usage, and wood Btu usage per household from: Energy Information Administration. Total Energy Consumption in U.S. Households by Type of Housing Unit, 2001  
<http://www.eia.doe.gov/emeu/recs/contents.html>.

CO<sub>2</sub>, SO<sub>2</sub>, and NO<sub>x</sub> power plant emission per kWh from: U.S. Environmental Protection Agency. U.S. Power Plant Emissions Total by Year  
[www.epa.gov/cleanenergy/egrid/samples.htm](http://www.epa.gov/cleanenergy/egrid/samples.htm).

CO emission per kWh assumes 1/3 of one percent of C emissions is CO based on: Energy Information Administration. 1994 Energy Use and Carbon Emissions: Non-OECD Countries DOE/EIA-0579.

PM<sub>10</sub> emission per kWh from: Layton, M. 2004. 2005 Electricity Environmental Performance Report: Electricity Generation and Air Emissions. California Energy Commission.  
[http://www.energy.ca.gov/2005\\_energy\\_policy/documents/2004-11-15\\_workshop/2004-11-15\\_03-A\\_LAYTON.PDF](http://www.energy.ca.gov/2005_energy_policy/documents/2004-11-15_workshop/2004-11-15_03-A_LAYTON.PDF)

CO<sub>2</sub>, NO<sub>x</sub>, SO<sub>2</sub>, PM<sub>10</sub>, and CO emission per Btu for natural gas, propane and butane (average used to represent LPG), Fuel #4 and #6 (average used to represent fuel oil and kerosene) from: Abraxas energy consulting, <http://www.abraxasenergy.com/emissions/>

CO<sub>2</sub> and fine particle emissions per Btu of wood from: Houck, J.E. Tiegs, P.E, McCrillis, R.C. Keithley, C. and Crouch, J. 1998. Air emissions from residential heating: the wood heating option put into environmental perspective. In: Proceedings of U.S. EPA and Air Waste Management Association Conference: Living in a Global Environment, V.1: 373-384.

CO, NO<sub>x</sub> and SO<sub>x</sub> emission per Btu based on total emissions and wood burning (tonnes) from: Residential Wood Burning Emissions in British Columbia, 2005.  
[http://www.env.bc.ca/air/airquality/pdfs/wood\\_emissions.pdf](http://www.env.bc.ca/air/airquality/pdfs/wood_emissions.pdf).

Emissions per dry tonne of wood converted to emissions per Btu based on average dry weight per cord of wood and average Btu per cord from: Heating with Wood I. Species characteristics and volumes. <http://ianrpubs.unl.edu/forestry/g881.htm>

22. Nowak, D.J. 1995. Trees pollute? A "TREE" explains it all. In: Proceedings of the 7th National Urban Forestry Conference. Washington, DC: American Forests. Pp. 28-30

23. Nowak, D.J. and J.F. Dwyer. 2007. Understanding the benefits and costs of urban forest ecosystems. In: Kuser, J. (ed.) Urban and Community Forestry in the Northeast. New York: Springer. Pp. 25-46.

24. Nowak, D.J. 2000. The interactions between urban forests and global climate change. In: Abdollahi, K.K., Z.H. Ning, and A. Appenning (Eds). Global Climate Change and the Urban Forest. Baton Rouge: GCRCC and Franklin Press. Pp. 31-44.

25. Nowak, D.J., D.E. Crane and J.C. Stevens. 2006. Air pollution removal by urban trees and shrubs in the United States. Urban Forestry and Urban Greening. 4:115-123

26. Nowak, D.J., R.E. Hoehn, D.E. Crane, J.C. Stevens, J.T. Walton, and J. Bond. 2008. A ground-based method of assessing urban forest structure and ecosystem services. Arboric. Urb. For. 34(6): 347-358

27. Hirabayashi, S., C. Kroll, and D. Nowak. (2011). Component-based development and sensitivity analyses of an air pollutant dry deposition model. *Environmental Modeling and Software* 26(6): 804-816.